

Lattice Electromagnetic Theory via Differential Forms

F. L. Teixeira, B. He

ElectroScience Laboratory and Department of Electrical Engineering
The Ohio State University

Fax (614) 292-7297, email: teixeira.5@osu.edu
<http://esl.eng.ohio-state.edu>

Summary - The theory of differential forms and exterior calculus are being employed for a rigorous representation of Maxwell's equations for finite-difference and finite-element time-domain simulations on irregular, unstructured lattices (grids). Differential forms and their discrete counterparts (co-chains) provide a natural bridge between the continuum and the discrete versions of the theory, and allow for a natural factorization of the equations into a metric part and a purely topological (i.e., invariant under homeomorphisms) part. The various sources of inconsistencies that commonly plague discretization schemes in irregular, unstructured grids can be more easily identified, classified, and avoided.


Motivation

- To obtain a discrete, coordinate-free representation of Maxwell's equations on a lattice which preserves the topological structure of the continuum theory at an exact level.
- Conservation laws and most theorems of the continuum are automatically satisfied, independently of the metric structure of the lattice (irregular or not).
- Metric approximations (interpolation, quadratures) are localized in a minimal number of operations.
- Classify the most common sources of inconsistency in the usual discretization schemes (FDTD, FEM, FVTD etc.).

Factorization of Maxwell's equations



Homeomorphic lattices with distinct metric structures.



Non-homeomorphic lattices

- Lattice equations invariant under homeomorphisms.
- Such invariance divides lattices into equivalence classes.
- All metric information is contained in the constitutive equations.

Maxwell's Equations in Forms Language

- Dynamic equations.

$$dE = i\omega B \quad (2a)$$


$$dH = -i\omega D + J_E \quad (2b)$$

$$dB = 0 \quad (2c)$$


$$dD = \rho_E \quad (2d)$$

- E, H : 1-forms, associated with edges.
- D, B : 2-forms, associated with faces.
- Constitutive equations are defined through Hodge operators \star , $D = \star E$, $B = \star H$ [isomorphism between space of k -forms and space of $(n - k)$ -forms].

Dual Lattices and Orientation



- E, B : ordinary forms with internal orientation (forces). H, D : twisted forms with external orientation (sources).



- Primary grid: internal. Dual grid: external. Different behavior under coordinate reflection.

Discrete Hodge Operators

- Procedural equivalents of $[\star]$ and $[\star_\rho]$ on vector calculus language are calculated through interpolation, quadratures. At this point, approximations are necessarily invoked.
- Interpolation of the dynamic variables on simplicial lattice (s_1^1, E) , (s_2^2, D) , (s_1^1, H) , (s_2^2, B) : Whitney forms.
- $k = 1$ case: $\Omega_{s_1^1} \xrightarrow{gE} \tau_{s_1^1} = \zeta_{i,0} \nabla \zeta_{i,1} - \zeta_{i,1} \nabla \zeta_{i,0}$
- Useful to avoid spurious modes in the FEM. Dual lattice: not simplicial anymore.

